

Carve™

For control of listed diseases on Turf

Active Ingredients:

Prothioconazole:41.00%

Other Ingredients:59.00%

Total:100.0%

Contains 4 lbs. Prothioconazole per gallon.

KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
IF SWALLOWED:	• Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
IF IN EYES:	• Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses if present, after the first 5 minutes, then continue rinsing. • Call a poison control center or doctor for treatment advice.
IF INHALED:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
Have the product container or label with you when calling a poison control center or doctor or going for treatment.	
Note to Physician: No specific antidote. Treat symptomatically.	
Emergency phone numbers	(800) 424-9300 CHEMTREC (transportation and spills) (800) 222-1222 Poison Control Center

Net Contents: 2.5 gallons (9.46 L)

EPA Reg. No. 60063-99

EPA Est. No. 60063-GA-1 (Lot No. begins with VL)
 EPA Est. No. 62171-MS-1 (Lot No. begins with OI)
 EPA Est. No. 70815-GA-1 (Lot no. begins with CB)
 EPA Est. No. 70989-MO-1 (Lot No. begins with ST)
 EPA Est. No. 72344-MO-4 (Lot no. begins with TR)
 EPA Est. No. 86555-MO-1 (Lot no. begins with AF)

Manufactured for:
Sipcam Agro USA, Inc.
2525 Meridian Parkway
Durham, NC 27713

Read the entire label carefully before opening the container.

See additional Precautionary Statements and Directions for Use inside booklet.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Caution. Harmful if swallowed or inhaled. Avoid breathing spray mist. Avoid contact with eyes or clothing. Causes moderate eye irritation. Remove and wash contaminated clothing before reuse. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PHYSICAL – CHEMICAL HAZARDS

DO NOT mix or allow to come in contact with oxidizing agents. Hazardous chemical reactions may occur.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

All handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Protective eyewear
- Chemical-resistant gloves made of any waterproof material

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to estuarine/marine invertebrates, and freshwater/estuaries/marine aquatic plants. For terrestrial uses: **DO NOT** apply directly to water, or to areas where surface water is present, or to

intertidal areas below the mean highwater mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Ground Water Advisory

Degradates of prothioconazole are known to leach through soil into groundwater under certain conditions as a result of label use. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

Surface Water Advisory

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having a high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of prothioconazole and degradates from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
Read the entire label before using this product.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

NON – AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are **NOT** within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. [Golf courses are not within the scope of the WPS.]

DO NOT enter or allow others to enter treated areas until sprays have dried.

PRODUCT INFORMATION

This product is a broad spectrum, preventative fungicide with systemic and curative properties. This product may be applied as a foliar spray and for soil application in alternating spray programs or in tank mixes with other registered crop protection products. All applications must be made according to the use directions that follow. Equipment must be properly calibrated before use.

SPRAY DRIFT

Aerial Applications:

- **DO NOT** release spray at a height greater than 10 ft. above the ground or crop canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select nozzle and pressure that deliver medium or coarser droplets in accordance with the most current version of the American Society of Agricultural & Biological Engineers Standard 641 (ASABE S641).
- **DO NOT** apply when wind speeds exceed 15 mph at the application site. If the windspeed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters
- Applicators must use 1/2 swath displacement upwind at the downwind edge of the field.
- **DO NOT** apply during temperature inversions.

Groundboom Applications:

- User must only apply with the release height recommended by the manufacturer, but no more than 4 feet above the ground or crop canopy.
- Applicators are required to use a Medium or coarser droplet size in accordance with the most current version of the American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- Do not apply when wind speeds exceed 15 miles per hour at the application site.
- Do not apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.
BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size – Aircraft

- Adjust Nozzles – Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

BOOM HEIGHT – Ground Boom

- For ground equipment, the boom should remain level with the target plants and have minimal bounce.

RELEASE HEIGHT – Aircraft

- Higher release heights increase the potential for spray drift.

SHIELDED SPRAYERS

- Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

TEMPERATURE AND HUMIDITY

- When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

TEMPERATURE INVERSIONS

- Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

WIND

- Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.
- Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

HANDHELD TECHNOLOGY APPLICATIONS

- Take precautions to minimize spray drift.

MIXING, LOADING, AND APPLYING

This product is intended to be diluted into water and then applied to crops by typical agricultural spraying techniques. Always apply this product in sufficient water to obtain thorough, uniform coverage of foliage and crop surfaces intended to be protected from disease. For spray volumes, see the individual crop charts. Both ground and aircraft methods of application are acceptable unless specific directions are given for a crop.

Slowly invert container several times to assure uniform mixture. Measure the required amount of this product and pour into the spray tank during filling. Keep agitator running when filling spray tank and during spray operations.

It is necessary to thoroughly apply the product in order to provide good disease control. **DO NOT** prepare more spray solution than is needed for application. Avoiding spray overlap will reduce the potential for crop injury.

Tank Mixing

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

When tank mixing this product with other pesticides, observe the more restrictive label limitations and precautions. **DO NOT** exceed any label dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing.

DO NOT combine this product in the sprayer tank with pesticides, surfactants or fertilizers, unless prior use has shown the combination to be physically compatible, non-injurious and effective under similar use conditions.

Compatibility Test for Mix Components: Before mixing components, always perform a compatibility jar test. To determine the physical compatibility, the specified proportions of products must be added into a suitable container of water in the following sequence:

1. Products packaged in water soluble packaging
2. Wettable powders, wettable granules (dry flowables)

3. Carve
4. Other liquid flowables
5. Emulsifiable concentrates
6. Water soluble liquids
7. Adjuvants

Using a quart jar, add the proportionate amounts of the products to 1 qt. of water. Add in the sequence indicated above. Always mix between component additions. After thoroughly mixing, let stand for at least 10-15 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

The crop safety of all potential tank-mixes, including additives and other pesticides, on all crops, has not been tested. Before applying any tank-mixture not specifically listed on this label, the safety to the target crop must be confirmed. To test for crop safety, apply the combination to a small area of the target crop in accordance with the label instructions to ensure that a phytotoxic response will not occur.

Applications through Sprinkler Irrigation Systems (Chemigation)

Apply this product only through center pivot, solid set, drip, linear, or moving wheel irrigation systems. **DO NOT** apply this product through any other type of irrigation system. Use only on crops specifically designated in the **Crop Use Directions**.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

DO NOT apply this product through irrigation systems connected to a public water system. 'Public water system' means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Controls for both irrigation water and pesticide injection systems must be functionally interlocked, so as to automatically terminate pesticide injection when the irrigation water pump motor stops. A person knowledgeable of the irrigation system and responsible for its operation must be present so as to discontinue pesticide injection and make necessary adjustments, if the need arises.

The irrigation water pipeline must be fitted with a functional, automatic, quick-closing check valve to prevent the flow of treated irrigation water back toward the water source. The pipeline must also be fitted with a vacuum relief valve and low pressure drain, located between the irrigation water pump and the check valve, to prevent back-siphoning of treated irrigation water into the water source.

Always inject this product into irrigation water after it discharges from the irrigation pump and after it passes through the check valve. Never inject pesticides into the intake line on the suction side of the pump.

Pesticide injection equipment must be fitted with a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump. Interlock this valve to the power system, so as to prevent fluid from being withdrawn from the chemical supply tank when the irrigation system is either automatically or manually turned off.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.

Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur. **DO NOT** apply when wind speed favors drift beyond the area intended for treatment.

This product may be used through two basic types of sprinkler irrigation systems as outlined in Sections A and B below. Determine which type of system is in place, then refer to the appropriate directions provided for each type.

A. Center Pivot, Motorized Lateral Move and Traveling Gun Irrigation Equipment

For injection of pesticides, these continuously moving systems must use a metering pump, such as a positive displacement injection pump of either diaphragm or piston type, constructed of materials that are compatible with pesticides, fitted with a system interlock, and capable of injection at pressures approximately 2 to 3 times those encountered within the irrigation water line. Venturi applicator units cannot be used on these systems.

Fill chemical supply tank of injection equipment with water. Operate system for one complete revolution or run across the field, measuring time required, amount of water injected, and acreage covered. Thoroughly mix the specified amount of this product for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run. Mixture in the chemical supply tank must be continuously agitated during the injection run. Shut off injection equipment after one revolution or run but continue to operate irrigation system until this product has been cleared from the last sprinkler head.

Use only with drive systems, which provide uniform water distribution. Do not use end guns when chemigating through center pivot systems because of non-uniform application.

B. Solid Set and Portable (Wheel Move, Side Roll, End Tow, or Hand Move) Irrigation Equipment

With stationary systems, an effectively designed in-line Venturi applicator unit is preferred which is constructed of materials that are compatible with pesticides; however, a positive-displacement pump can also be used.

Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a 30 – 45 minute period. Mix desired amount of this product for acreage to be covered with water so that the total mixture of this plus water in the injection tank is equal to the quantity of water used during calibration and operate entire system at normal pressures specified by the manufacturer of injection equipment used for the amount of time established during calibration. No agitation should be required. This product can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop injection equipment after treatment is completed and continue to operate irrigation system until this product has been cleared from the last sprinkler head.

INTEGRATED PEST / DISEASE MANAGEMENT

This product provides excellent control of fungal diseases when used according to label directions for control of a broad spectrum of plant diseases. This product is for use in programs that are compatible with the principals of Integrated Pest Management (IPM), including the use of disease resistant crop varieties, cultural practices, pest scouting, and disease forecasting systems which reduce unnecessary applications of pesticides.

RESISTANCE MANAGEMENT

For resistance management, this product contains prothioconazole, a DMI Group 3 Fungicide. Any fungal population may contain individuals naturally resistant to this product and other Group 3

fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Group 3 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological, and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Sipcam Agro at 919-226-1195. You can also contact your pesticide distributor or university extension specialist to report resistance.

TURF USE DIRECTIONS

This product controls a range of turf diseases. Refer to the table below for disease, rate structure, and specific use information. Apply this product prior to turf disease development or in the early stages of turf disease development.

APPLICATION DIRECTIONS

Apply this product through ground application equipment with boom below and behind the operator to target soil or foliar turfgrass diseases. See 'Turf Use Directions' for application rates and use information. Turfgrass injury and/or inferior disease control may result upon failure to follow label directions.

Ground Application Equipment and Nozzles

Before use, ensure equipment is properly calibrated. Use nozzles that are accurate, and provide uniform coverage to the turfgrass foliage and/or the soil surface. Use the same size nozzles, uniformly distributed across the boom, and operate nozzles within the spray pressure directions suggested by the nozzle manufacturer.

Spray Equipment Preparation and Mixing Instructions

Prepare no more spray mixture than is necessary for the immediate application. Start with clean spray equipment before using this product. While preparing the mixture, and throughout the entire spray operation, maintain agitation. **DO NOT** let the spray mixture stand overnight in the spray tank. Following use, flush the spray equipment thoroughly. Rinsate may be applied to the previously treated areas, or disposed of according to local regulations.

Mixing Instructions for Carve alone

- Add 1/4 of the required amount of water to the mix tank.
- With the agitator running, add Carve to the mix tank.
- Continue agitation while adding the remainder of the water.
- Begin application after the product has completely and uniformly dispersed into the mix water.
- Maintain agitation until all of the mixture has been applied.

Application Volume

Apply this product in a minimum of 40 gallons of spray solution per acre. Complete coverage and uniform applications are essential for most effective results, especially when lower spray volumes are applied. If necessary, increase spray volume for uniform and complete coverage.

USE RESTRICTIONS

- For use only on golf course turfgrass (greens, tees, fairways, and roughs).
- **NOT FOR RESIDENTIAL USE. DO NOT** use on home lawns and turf sites associated with apartment buildings, daycare centers, playgrounds, playfields, recreational park, athletic fields, schools, campgrounds, churches, and theme parks.
- **DO NOT** use clippings for animal feed.
- **DO NOT** apply by aerial application.
- **DO NOT** apply through any type of irrigation equipment (chemigation).
- For ground application equipment only. **DO NOT** use with handheld application equipment, including mechanically pressurized spray gun, backpack or tank pressurized spray gun, spray hawk or handheld boom applicators.

USE PRECAUTIONS

- After application, allow foliage to dry prior to mowing.
- Use of skirted spray booms will further reduce spray drift.

Disease	Application Rate ¹	Interval between Applications (days) ²	Use Information
Anthrachnose (<i>Colletotrichum graminicola</i>)	8.5 (fl.oz. /A)	14 – 21	
Brown Patch [Rhizoctonia Blight] (<i>Rhizoctonia solani</i>)	8.5 (fl.oz. /A)	14 – 21	For best results, apply preventatively before conditions favorable for infection occur.
Brown Ring (Waitea) Patch (<i>Waitea circinata</i> var. <i>circinata</i>)	8.5 (fl.oz. /A)	14 – 21	For best results, apply preventatively before conditions favorable for infection occur.
Dollar Spot (<i>Sclerotinia homoeocarpa</i> syn. <i>Clariireedia homoeocarpa</i>)	8.5 (fl.oz. /A)	14 – 21	
Fairy Ring (caused by Basidiomycete fungi)	8.5 (fl.oz. /A)	14 – 28	Before the spray dries, adequately water-in to move the fungicide into the thatch/soil where the fungus is active.
Gray Leaf Spot (<i>Pyricularia grisea</i>)	8.5 (fl.oz. /A)	14 – 21	
Gray Snow Mold [Typhula Blight] [(<i>Typhula incarnata</i>)] [(<i>Typhula incarnata</i> , T. <i>ishikariensis</i>)] [*] Pink Snow Mold (<i>Microdochium nivale</i>)	8.5 (fl.oz. /A)	14 – 28	Make 1-2 preventative applications before conditions favorable for infection occur. Do not apply over snow cover.
Large Patch [Zoysia Patch] (<i>Rhizoctonia solani</i>)	8.5 (fl.oz. /A)	14 – 28	Lightly water-in applications to move fungicide into crown area and root zone.
Microdochium Patch (<i>Microdochium nivale</i>)	8.5 (fl.oz. /A)	14 – 21	Begin preventative applications when conditions are conducive to disease development and without lasting snowfall.
Mini Ring (<i>Rhizoctonia zeae</i>)	8.5 (fl.oz. /A)	14 – 21	Apply preventatively before conditions favorable for infection occur.

Spring Dead Spot (<i>Ophiosphaerella korrae</i> , <i>O. herpotricha</i> , <i>Leptosphaeria korrea</i> , <i>L. namari</i>)	8.5 (fl.oz. /A)	14 – 28	Lightly water-in applications to move fungicide into crown area and root zone.
Summer Patch (<i>Magnaporthe poae</i>)	8.5 (fl.oz. /A)	14 – 28	Use adequate spray-volume or water-in applications to crowns and upper roots for maximum effectiveness. If there is prior disease history, use the higher rate and shorter interval.
Take-all Patch (<i>Gaeumannomyces ramms</i> var. <i>avenae</i>)	8.5 (fl.oz. /A)	14 – 21	Treat preventatively for the disease. Water-in applications into the upper root zone.
Take-all Root Rot (<i>Gaeumannomyces graminis</i> var. <i>graminis</i>)	8.5 (fl.oz. /A)	14 – 28	Apply fungicide in adequate water volume or water-in applications to upper root zone.
¹ Rate of 8.5 fl.oz. product/A equates to 0.196 fl.oz. product/1000 sq.ft. or 0.267 lbs.AI/A. ² Use the shorter specified application interval and/or the higher specified rate when prolonged favorable turf disease conditions exist.			
RESTRICTIONS • Maximum Single Rate: DO NOT exceed 8.5 fl.oz. product/A per application (0.196 fl.oz. product/1,000 ft²), equivalent to 0.267 lbs a.i./A. • Yearly Maximum Rate: DO NOT exceed 25.6 fl.oz. product/A per year (0.588 fl.oz. product/1,000 ft²/year), equivalent to 0.801 lbs a.i./A/year. • DO NOT exceed 3 applications per year using the maximum single application rate.			

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

STORAGE

Store in original containers only. Keep container closed when not in use. Do not store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to the label.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative of the nearest EPA Regional Office for guidance.

CONTAINER HANDLING [less than or equal to 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill.

CONTAINER HANDLING [greater than 5 gallons]

Refillable container. Refill this container with pesticide only. Do not reuse the container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

CONTAINER HANDLING [greater than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container or pressure rinse promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full of water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by other procedures allowed by state and local authorities.

CONTAINER IS NOT SAFE FOR FOOD, FEED OR DRINKING WATER.

WARRANTY AND LIMITATION OF DAMAGES

CONDITIONS OF SALE: To the extent consistent with applicable law, Sipcam Agro USA, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Sipcam Agro USA, Inc. SIPCAM AGRO USA, INC. DISCLAIMS ALL OTHER WARRANTIES, EXPRESS OR IMPLIED. To the extent consistent with applicable law, SIPCAM AGRO USA, INC. SHALL NOT BE LIABLE FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, AND SIPCAM AGRO USA, INC.'S SOLE LIABILITY AND BUYER'S AND USER'S EXCLUSIVE REMEDY SHALL BE LIMITED TO THE REFUND OF THE PURCHASE PRICE. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER AND USER ACKNOWLEDGE AND ASSUME ALL RISKS AND LIABILITY RESULTING FROM HANDLING, STORAGE AND USE OF THIS PRODUCT. SIPCAM AGRO USA, INC. DOES NOT AUTHORIZE ANY AGENT OR REPRESENTATIVE TO MAKE ANY OTHER WARRANTY, GUARANTEE OR REPRESENTATION CONCERNING THIS PRODUCT.