

Active Ingredient: Water-washed Kaolin 100%



Cocoon® is a multi-functional solution to protecting your crop investment. Cocoon® promotes higher yields and the harvest of more marketable fruits and vegetables. Manufactured with a safened formulation, Cocoon®'s water-washed kaolin process protects fruit against direct sunburn and heat stress damage when applied. Cocoon® can reduce sunburn losses by up to 50% while keeping plant canopies cooler, which leads to more efficient photosynthesis and higher yields in high light and heat environments.

- High quality active ingredient forms a reflective barrier to protect your crop
- Prevents damage from high temperatures
- Easy-to-use in conventional sprayers, the nonabrasive formula can be mixed with a surfactant for optimum coverage and adhesion
- Can be tank-mixed with most crop protection products and nutrients

- Stress protection allows for higher crop yield
- Healthy plant growth increases even during stressful events
- Allows for additional surfactant use which can improve coverage on plant surface
- Fits both conventional and organic production programs



Used On

For use on pome fruit, stone fruit, citrus, tree nuts, berries, grapes, fruiting vegetables, legumes, bulb vegetables, cucurbits, roots and tubers, brassica vegetables, leafy vegetables, cotton, miscellaneous tree and vine crops, nursery and ornamental plants and Christmas trees.

To learn more about Cocoon® and our entire portfolio of agricultural solutions, talk to your Sipcam Agro USA representative or visit www.sipcam.com. Always read and follow label directions.

Sipcam Agro USA

2525 Meridian Parkway Suite 100 | Durham, NC | sipcam.com | (<u>877) 898-9514</u>

©2025 Sipcam Agro USA, Inc. All rights reserved.





Technical Information

Preventative Care

Sunburn damage on fruit, vegetable and nut crops can put a blemish on harvest profits. Apply Cocoon® before solar radiation does its damage. Cocoon® places a high quality particle film protective barrier of kaolin clay around food, nursery and greenhouse crops to reduce heat stress.





Protective cooling, reflective white barrier







Untreated

How to Use

- Make the initial application at the highest recommended rate. Mix with water and apply Cocoon® to fruit, foliage, limb and trunk surfaces before periods of high temperatures and/or intense sunlight
- Cocoon® forms a dry, white reflective barrier
- A second application of Cocoon® can be made after 7 to 10 days for adequate coverage; subsequent applications may be made at 7 to 21-day intervals
- Easy-to-use and can be applied by ground or air in sufficient water for thorough coverage
- The addition of a surfactant may aid in improving coverage on plant surface
- Cocoon® can be tank-mixed with most crop protection products and nutrients

Mixing Instructions

- For concentrate sprays, use no less than 10 U.S. gal/acre
- For ground application, 100-200 U.S. gal/acre of water is suggested in order to achieve thorough coverage on plant surfaces; avoid excessive runoff
- Use the rates suggested for each crop listed; normal application rates are 25-50 lbs/acre
- See label for complete crop recommendation

To learn more about Cocoon® and our entire portfolio of agricultural solutions, talk to your Sipcam Agro USA representative or visit **www.sipcam.com**Always read and follow label directions.

Sipcam Agro USA

2525 Meridian Parkway Suite 100 | Durham, NC | sipcam.com | (877) 898-9514



