



Active ingredients: cymoxanil, chlorothalonil

# THE SMART CHOICE FOR LATE BLIGHT CONTROL



Late blight can do more than negatively impact yields or returns. Without proper management, a late blight outbreak can decimate an entire crop. While there are many steps in preventing and managing late blight, the use of foliar fungicides, like Cymbol™ Advance, on a regular and continuing schedule is the only way to manage the disease in the field. Cymbol Advance combines the unique resistance management benefits of chlorothalonil with the excellent preventative and curative activity of cymoxanil. The combination of these powerful modes of action give potato growers both contact and locally systemic activity, resulting in full-season, multi-level protection for their potato crop.





## DISCOVER DISEASE PROTECTION THAT DOESN'T QUIT

- » Strong locally systemic action to fight late blight from within the plant
- » Superior sporulation control reduces future cycles of disease by inhibiting sporulation
- » Combination of multi-site, broad-spectrum chlorothalonil plus the preventative, curative and antispore activity of cymoxanil offers exceptional resistance management
- » Economical disease management combined with the highest disease control performance results in proven ROI

## FIELD PROVEN

Cymbol Advance puts potato growers in the driver's seat when it comes to interval frequency. The product's flexibility allows growers to make decisions regarding applications based on the unique circumstances in their field for any particular growing season. Fields treated with the Cymbol Advance program are proven to provide comparable disease control and yields to those treated with the Grower Standard Program.

### HUBBARD AGRICULTURAL SCIENCE POTATO FUNGICIDE TRIAL



## RECOMMENDED USE RATES

2 pints/A (0.09 to 0.1275 lb. a.i./A)

- » Cymbol Advance also controls downy mildew and late blight diseases on tomatoes, cucurbits and fruiting vegetables. See label for complete list of crops, diseases and rates.
- » Timing: Every 14 days or earlier before harvest
- » Buffer tank solution to a pH of 7 or below prior to the addition of Cymbol Advance

## PACKAGING

2 x 2.5 gallons per case, 260 gallon tote

## FRAC CODE

27 (cymoxanil), M5 (chlorothalonil)

Always read and follow label directions. ©2020 Sipcam Agro USA, Inc. The Sipcam and Cymbol Advance logos are registered trademarks of Sipcam Agro USA, Inc. Echo 720 is registered product of Sipcam Agro USA, Inc. Reason and Previcur are registered products of Bayer.

## REASONS FOR EARLY SEASON APPLICATION

- » Secures control of disease source, attacking the first disease cycle
- » Manages and contains disease before it spreads to other plants
- » Protects and covers stems from disease before row closure
- » Stops unseen infection up to three days after initial infection
- » Offers the most economic approach to disease management, delivering the best ROI and peace of mind

**Real People. Real Solutions.** To learn more about Cymbol Advance or our entire portfolio, talk to your Sipcam Agro distributor; call 1-877-898-9514 or visit [sipcamagro.com](http://sipcamagro.com).

[sipcam.com](http://sipcam.com)

