

# RESTRICTED USE PESTICIDE

## DUE TO GROUND AND SURFACE WATER CONCERNS

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION. THIS PRODUCT IS A RESTRICTED-USE HERBICIDE DUE TO GROUND AND SURFACE WATER CONCERNS. USERS MUST READ AND FOLLOW ALL PRECAUTIONARY STATEMENTS AND INSTRUCTIONS FOR USE IN ORDER TO MINIMIZE POTENTIAL FOR ATRAZINE TO REACH GROUND AND SURFACE WATER

# Stalwart® 3W

Metolachlor	Group	15	Herbicide
Mesotrione	Group	27	Herbicide
Atrazine	Group	5	Herbicide

For pre-emergence and post-emergence control of grass and broadleaf weeds in Corn (field, seed, silage, sweet, and yellow popcorn), and Grain Sorghum

### ACTIVE INGREDIENTS:

Metolachlor	19.00%
Atrazine	18.60%
Mesotrione	2.44%

### OTHER INGREDIENTS:

	59.96%
--	--------

TOTAL: 100.00%

Equivalent to 1.70 lbs.ai./gal. metolachlor; 0.218 lbs.ai./gal. mesotrione; and 1.66 lbs.ai./gal. atrazine.

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

### FIRST AID

#### IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

#### IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

#### IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

#### IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Emergency Phone Numbers

Poison Control: 1-800-222-1222  
CHEMTREC: 1-800-424-9300 (transportation and spills)

See additional Precautionary Statements and Directions for Use inside the booklet.

Manufactured for:  
Sipcam Agro USA, Inc.  
2525 Meridian Parkway  
Durham, NC 27713

For Spanish translation, see inside the booklet. (Para la traducción al español, consulte el interior del folleto.)

EPA Reg. No.: 60063-64

EPA Est. No.: 60063-GA-1 (Lot number begins with VL)

EPA Est. No.: 62171-MS-1 (Lot number begins with OI)

EPA Est. No.: 70989-MO-1 (Lot number begins with ST)

EPA Est. No.: 72344-MO-4 (Lot number begins with TR)

EPA 20230120 (9/25)



**SIPCAM AGRO**  
USA, INC.

Not for sale, use, or distribution in Nassau County or Suffolk County, New York.

**READ THE LABEL  
CAREFULLY  
BEFORE OPENING  
THE CONTAINER**

HERBICIDE

## PESTICIDA DE USO RESTRINGIDO

### DEBIDO A CUESTIONES RELACIONADAS CON LAS AGUAS SUBTERRÁNEAS Y SUPERFICIALES.

PARA LA VENTA AL POR MENOR Y EL USO POR APLICADORES CERTIFICADOS O PERSONAS QUE ESTÉN BAJO SU DIRECTA SUPERVISIÓN Y ÚNICAMENTE PARA LOS USOS DETALLADOS EN LA CERTIFICACIÓN DEL APLICADOR CERTIFICADO. ESTE PRODUCTO ES UN HERBICIDA DE USO RESTRINGIDO DEBIDO A CUESTIONES RELACIONADAS CON LAS AGUAS SUBTERRÁNEAS Y SUPERFICIALES. LOS USUARIOS DEBEN LEER Y SEGUIR TODAS LAS DECLARACIONES PREVENTIVAS E INSTRUCCIONES DE USO PARA REDUCIR AL MÍNIMO LA POSIBILIDAD DE QUE LA ATRAZINA CONTAMINE LAS AGUAS SUBTERRÁNEAS Y SUPERFICIALES.

# Stalwart 3W

## KEEP OUT OF REACH OF CHILDREN

### MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

#### CAUTION

#### PRECAUCIÓN

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

#### PRIMEROS AUXILIOS

<b>SI ENTRA EN CONTACTO CON LOS OJOS:</b>	<ul style="list-style-type: none"><li>• Mantenga los ojos abiertos y enjuáguelos lenta y cuidadosamente con agua, durante 15 a 20 minutos.</li><li>• Si utiliza lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando.</li><li>• Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.</li></ul>
<b>SI SE INGIERE:</b>	<ul style="list-style-type: none"><li>• Llame de inmediato a un centro de control de envenenamientos o a un médico para consejo de tratamiento.</li><li>• Si la persona puede tragar, haga que beba un vaso de agua lentamente.</li><li>• No induzca al vómito, a menos que así se lo indique un centro de control de envenenamientos o un médico.</li><li>• No administre nada por boca a una persona que haya perdido el conocimiento.</li></ul>
<b>SI CAE EN LA PIEL O LA ROPA:</b>	<ul style="list-style-type: none"><li>• Quítense la ropa contaminada.</li><li>• Enjuague la piel inmediatamente con bastante agua por 15-20 minutos.</li><li>• Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.</li></ul>
<b>SI ES INHALADO:</b>	<ul style="list-style-type: none"><li>• Traslade a la persona al aire fresco.</li><li>• Si la persona no está respirando llame al 911 o a una ambulancia, luego dé respiración artificial, preferiblemente de boca a boca, si es posible.</li><li>• Llame al centro de control de envenenamientos o a un médico para consejo de tratamiento.</li></ul>
Cuando llame a un centro de control de envenenamientos, o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto.	
El Número de Emergencia	Para emergencias médicas, llame al centro de control de envenenamientos a (800)-222-1222. Para emergencias derrames de emergencia, llame CHEMTREC a (800)-424-9300.

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS & DOMESTIC ANIMALS

**CAUTION.** Causes moderate eye irritation. Harmful if swallowed. Avoid contact with skin, eyes, or clothing. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

## DECLARACIONES PREVENTIVAS

### RIESGOS PARA LOS SERES HUMANOS Y ANIMALES DOMÉSTICO

**PRECAUCIÓN.** Causa irritación moderada en los ojos. Dañino si se ingiere. No permita que caiga en los ojos, la piel o la ropa. El contacto prolongado y frecuente con la piel puede causar reacciones alérgicas en algunos individuos. Lávese con abundante agua y jabón luego de manejar (pesticidas) y antes de comer, beber, masticar chicle o tabaco, o ir al baño.

## PERSONAL PROTECTIVE EQUIPMENT (PPE)

### All Mixers, Loaders, Applicators, Flaggers, and other handlers must wear:

- Coveralls over short-sleeved shirt and short pants.
- Waterproof gloves
- Chemical-resistant footwear plus socks.
- Chemical-resistant apron, when mixing/loading, cleaning up spills, or cleaning equipment, or otherwise exposed to the concentrate.
- Chemical-resistant headgear for overhead exposure.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

### EQUIPO DE PROTECCIÓN PERSONAL (PPE, por sus siglas en inglés)

#### Los mezcladores, los cargadores, los aplicadores, los banderilleros y otros manipuladores de pesticidas (plaguicidas) deben usar:

- Overol (mameluco) encima de una camisa de manga corta y pantalones cortos.
- Guantes impermeables.
- Calcetines y zapatos resistentes a productos químicos
- Delantal resistente a productos químicos al mezclar o cargar, limpiar derrames, limpiar equipos o estar expuesto de cualquier otra manera al concentrado.
- Equipo de protección para la cabeza resistente a productos químicos en caso de exposición por encima de la cabeza.

Siga las instrucciones del fabricante para la limpieza o el mantenimiento del equipo de protección personal (PPE, por sus siglas en inglés). En caso de no existir dichas instrucciones de lavado, utilice detergente y agua caliente. Mantenga y lave el PPE separado de otra ropa para lavar. Deseche la ropa y otros materiales absorbentes que se hayan empapado o contaminado con el concentrado de este producto. No los reutilice.

### ENGINEERING CONTROL STATEMENTS

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(5)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### DECLARACIONES DE CONTROL TÉCNICO

Si los manipuladores emplean sistemas cerrados o cabinas cerradas de forma tal que se cumpla con los requisitos del Estándar para la Protección del Trabajador Agrícola (WPS, por sus siglas en inglés) para pesticidas agrícolas [40 CFR 170.240(d)(5)], los requisitos de PPE para los manipuladores pueden ser reducidos o modificados conforme a lo que establece el WPS.

### USER SAFETY RECOMMENDATIONS

#### Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### RECOMENDACIONES DE SEGURIDAD PARA EL USUARIO

#### Los usuarios deben cumplir con lo siguiente:

- Lavarse las manos antes de comer, beber, mascar chicle, consumir tabaco o ir al baño.
- Quitarse la ropa o el PPE inmediatamente si el pesticida penetra estas prendas. Luego, lavarse cuidadosamente y ponerse ropa limpia. Si el pesticida cae en la piel, lavarse inmediatamente con agua y jabón.
- Quitarse el PPE inmediatamente después de manipular este producto. Lavar el exterior de los guantes antes de quitárselos. A continuación, lavarse bien y cambiarse de ropa lo antes posible.

### ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinseate.

### Non-Target Organism Advisory Statement

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

### RIESGOS AMBIENTALES

Este producto es tóxico para los invertebrados acuáticos. No aplique directamente a cuerpos de agua, en áreas donde haya aguas superficiales, o en áreas intermareales que se encuentren bajo el nivel de altura máxima promedio del agua. No lo aplique cuando las condiciones meteorológicas favorezcan que el producto se aleje de las zonas tratadas. La escorrentía y la deriva del producto de las áreas tratadas puede suponer un peligro para los organismos acuáticos en las zonas circundantes. No contamine cuerpos de agua cuando disponga de las aguas de lavado de equipo o aguas de enjuague.

### Declaración consultiva sobre organismos no objetivo

Este producto es tóxico para las plantas y puede afectar negativamente al forraje y al hábitat de organismos no objetivo, incluidos los polinizadores, en zonas adyacentes al lugar tratado. Proteja el forraje y el hábitat de los organismos no objetivo siguiendo las instrucciones de la etiqueta destinadas a minimizar la deriva de la pulverización.

## GROUND WATER ADVISORY

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply atrazine to sand and loamy sand soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e., well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

Metolachlor and mesotrione are known to leach through soil into groundwater under certain conditions as a result of label use. These chemicals may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

### Surface Water Advisory

This product may impact surface water quality due to runoff of rainwater. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of metolachlor or mesotrione from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

### AVISO SOBRE AGUAS SUBTERRÁNEAS

La atrazina puede desplazarse (infiltrarse o lixiviarse) a través del suelo y llegar a las aguas subterráneas, que pueden utilizarse como agua potable. Se ha encontrado atrazina en aguas subterráneas. Se recomienda a los usuarios no aplicar la atrazina en suelos arenosos y franco-arenosos donde el nivel freático (las aguas subterráneas) esté cerca de la superficie y donde estos suelos sean muy permeables, es decir, drenen bien. Sus agencias agrícolas locales pueden proporcionarle más información sobre el tipo de suelo en su área y la ubicación de las aguas subterráneas.

Se sabe que el metolachlor y mesotrione pueden lixiviar a través del suelo y llegar a las aguas subterráneas bajo ciertas condiciones como resultado del uso conforme a la etiqueta. Estos químicos se puede filtrar a aguas subterráneas si es utilizado en áreas donde el suelo es permeable, especialmente en donde la tabla de agua está cercana a la superficie.

### Aviso sobre aguas superficiales

Este producto puede impactar la Calidad de cuerpos de agua superficiales debido a escorrentías de aguas pluviales. Esto es especialmente notable en suelos con poco drenaje y suelos con una tabla de agua que está cercana a la superficie. Este producto está clasificado como de alto potencial para alcanzar el agua superficial a través de la escorrentía durante varias semanas o meses después de su aplicación. Una franja de vegetación nivelada y bien mantenida entre las áreas donde se aplica este producto y las fuentes de agua superficial, como estanques, arroyos y manantiales, reducirá la carga potencial del metolachlor o mesotrione en el agua de escorrentía y los sedimentos. La escorrentía de este producto se reducirá si se evitan las aplicaciones cuando va a llover o se va a regar dentro de las 48 horas siguientes. La aplicación de prácticas efectivas de control de la erosión reducirá el potencial de este producto para alcanzar los sedimentos acuáticos a través de la escorrentía.

### RESTRICTIONS

This product must be used in a manner that will prevent back siphoning into wells and prevent spills.

Dispose of excess pesticide, spray mixtures or rinsates properly.

Mixing equipment must have check valves or anti-siphoning devices in use.

This product must not be mixed or loaded within 50 feet of perennial or intermittent streams and rivers, natural or impounded lakes and reservoirs. This product must not be applied within 66 feet of points where field surface water runoff enters perennial or intermittent streams and rivers or within 200 feet of natural or impounded lakes and reservoirs. If this product is applied to highly erodible land, the 66 foot buffer or setback from runoff entry points must be planted to crop, or seeded with grass or other suitable crop.

This product must not be mixed or loaded, or used within 50 feet of all wells, including abandoned wells, drainage wells, and sink holes. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 ft. of any well are prohibited, unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rain water that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site.

Additional State-imposed requirements regarding well-head setbacks and operational area containment must be observed.

Tile-Outletted Terraced Fields Containing Standpipes: One of the following restrictions must be used when applying atrazine to tile-outletted terraced fields containing standpipes:

- Do not apply within 66 ft. of standpipes in tile-outletted terraced fields.
- Do this product to the entire tile-outletted terraced field and immediately incorporate it to a depth of 2 to 3 inches in the entire field.
- Apply this product to the entire tile-outletted field under a no-till practice only when a high crop residue management practice is practiced. High crop residue management is described as a crop management practice where little or no crop residue is removed from the field during and after crop harvest.

## RESTRICCIÓN:

Este producto debe utilizarse de manera que se evite el reflujó a los pozos y se eviten los vertidos. Elimine adecuadamente el exceso de plaguicida, las mezclas de pulverización o los enjuagues.

Los equipos de mezcla deben disponer de válvulas de retención o dispositivos antisifón.

El producto no debe mezclarse ni cargarse a menos de 50 pies (15 metros) de arroyos y ríos intermitentes, lagos naturales o embalsados y embalses. El producto no debe aplicarse a menos de 66 pies (20 metros) de los puntos donde la escorrentía superficial del campo ingresa a arroyos y ríos perennes o intermitentes, ni a menos de 200 pies (61 metros) de lagos naturales o embalsados y embalses. Si este producto se aplica en terrenos altamente erosionables, la franja de protección o retroceso de 66 pies (20 metros) desde los puntos de entrada de la escorrentía debe plantarse con cultivos, o sembrarse con césped u otro cultivo adecuado.

El producto no debe mezclarse ni cargarse, ni usarse a menos de 50 pies (15 metros) de todos los pozos, incluidos los pozos abandonados, pozos de drenaje y sumideros. Se prohíben las operaciones que impliquen la mezcla, carga, enjuague o lavado de este producto en equipos o recipientes de manejo o aplicación de pesticidas o desde ellos a menos de 50 pies (15 metros) de cualquier pozo, a menos que se realicen sobre una plataforma impermeable construida para soportar el peso de la carga más pesada que pueda colocarse o trasladarse sobre la plataforma. Dicha plataforma deberá ser diseñada y mantenida para contener cualquier derrame de producto o fuga de equipo, agua de enjuague o lavado de contenedores o equipos, y agua de lluvia que pueda caer sobre la plataforma. No se debe permitir que el agua superficial fluya sobre la plataforma o desde ella, lo que significa que la plataforma debe ser autónoma. La plataforma deberá estar inclinada para facilitar la retirada del material. Una plataforma sin techo deberá tener una capacidad suficiente para contener, como mínimo, el 110 % de la capacidad del recipiente de pesticida o equipo de aplicación más grande sobre la plataforma. Una plataforma cubierta por un techo de tamaño suficiente para evitar que las precipitaciones entren en contacto con la plataforma deberá tener una capacidad mínima de contención del 100 % de la capacidad del recipiente de pesticida o equipo de aplicación más grande sobre la plataforma. Las capacidades de contención descritas anteriormente deberán mantenerse en todo momento. Las capacidades mínimas de contención especificadas anteriormente no se aplican a los vehículos que entregan plaguicidas en los lugares de mezcla/carga.

Deben cumplirse los requisitos adicionales impuestos por el estado en relación con los retranqueos de las cabezas de pozo y la contención de la zona de operaciones.

**Para la aplicación de atrazina en campos de cultivo en terrazas con baldosas que contengan tubos verticales, se debe utilizar una de las siguientes restricciones:**

- NO aplique este producto a menos de 66 pies (20 metros) de distancia de tuberías verticales en campos aterrizados con salida de baldosas.
- Aplique este producto a todo el campo aterrizado con baldosas e incorpóreló inmediatamente a una profundidad de 2 a 3 pulgadas (5 a 8 centímetros) en todo el campo.
- Aplique este producto a todo el campo aterrizado con salida de tejas bajo una práctica de labranza cero sólo cuando se realice una práctica de gestión de altos residuos de cultivos. La gestión de altos residuos de cultivos se describe como una práctica de gestión de cultivos en la que se retira poco o ningún residuo de cultivo del campo durante y después de la cosecha.

## PHYSICAL AND CHEMICAL HAZARDS

**DO NOT** use or store near heat or open flame. **DO NOT** mix or allow to come in contact with oxidizing agents, as a hazardous chemical reaction may occur.

## QUÍMICA FÍSICA

**NO** utilice ni almacene cerca del calor o llamas abiertas.

**NO** mezcle ni permita que entre en contacto con agentes oxidantes, ya que podría ocurrir una reacción química peligrosa.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

El uso de este producto de forma contraria a lo indicado en su etiqueta constituye una infracción de la ley federal. Leer toda la etiqueta antes de usar este producto.

## ENDANGERED SPECIES

It is a Federal offense to use any pesticide in a manner that results in an unauthorized "take" (e.g., kill or otherwise harm) of an endangered species and certain threatened species, under the Endangered Species Act section 9. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins, consult <http://www.epa.gov/esppl>, call 1-844-447-3813, or email [ESPP@epa.gov](mailto:ESPP@epa.gov). You must use the Bulletin valid for the month in which you will apply the product.

**ANY USE OF THIS PRODUCT IN AN AREA WHERE USE IS PROHIBITED IS A VIOLATION OF FEDERAL LAW.** Before using this product, you must consult the Atrazine Watershed Information Center (AWIC) to determine whether the use of this product is prohibited in your watershed. AWIC can be accessed through [www.atrazine-watershed.info](http://www.atrazine-watershed.info) or 1-866-365-3014. If use of this product is prohibited in your watershed, you may return this product to your point of purchase or contact Sipcam Agro USA, Inc. for a refund.

Use of this product in certain portions of California, Oregon, and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et al. v. EPA, C01-0132C, (W.D. WA). For further information, please refer to <https://www.epa.gov/endangered-species/endangered-speciescase-washington-toxics-coalition-v-epa>.

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**NOT** for use in Hawaii, Alaska, and the U.S. territories (Puerto Rico, Guam, American Samoa, the U.S. Virgin Islands, and the North Mariana Islands).

Use on roadsides; Conservation Reserve Program (CRP) land; conifers, including Christmas Tree plantings; timber; forestry; and, Miscanthus and other perennial bioenergy crops is **prohibited**.

**DO NOT** apply atrazine and propazine products to the same sorghum acre.

Failure to follow the **DIRECTIONS FOR USE, RESTRICTIONS** and **PRECAUTIONS** on this label may result in reduced weed control, adverse crop response, or illegal crop residues.

**NOTE: Not for sale, distribution or use in Nassau or Suffolk Counties in New York.**

#### Reporting Ecological Incidents:

To report ecological incidents, including mortality, injury, or harm to plants and animals, call Sipcam Agro USA at 1-919-226-1288.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours. Exception:** If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated. The following PPE is required for early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water:

- Coveralls over short-sleeved shirt and short pants
- Chemical-resistant footwear plus socks
- Waterproof gloves
- Chemical-resistant headgear for overhead exposure.

#### REQUISITOS DE USO AGRÍCOLA

Utilice este producto solo de acuerdo con su etiqueta y el Estándar para la Protección del Trabajador Agrícola (WPS), 40 CFR Parte 170. Esta ley contiene los requisitos para la protección de los trabajadores agrícolas en granjas, bosques, viveros e invernaderos, y para las personas que manipulan pesticidas agrícolas. Contiene requisitos para la capacitación, descontaminación, notificación y asistencia de emergencia. También contiene instrucciones específicas y excepciones relacionadas con las indicaciones en esta etiqueta acerca del equipo de protección personal (PPE) y el intervalo de ingreso restringido. Los requisitos en esta sección de la etiqueta (requisitos para uso agrícola) aplican únicamente a los usos de este producto que están cubiertos por el Estándar para la Protección del Trabajador Agrícola. **No ingrese o permita el ingreso de trabajadores a las áreas tratadas durante el intervalo de acceso restringido (REI, por sus siglas en inglés) de 24 horas. Excepción:** Si el producto se inyecta o incorpora a la tierra, el Estándar para la Protección del Trabajador Agrícola habilita, en determinadas circunstancias, que los trabajadores ingresen al área tratada siempre que no entren en contacto con nada que haya sido tratado. Se requiere el siguiente PPE para el acceso anticipado a áreas tratadas de acuerdo con el Estándar para la Protección del Trabajador Agrícola, y que involucre contacto con material tratado, como plantas, tierra o agua:

- Overol encima de una camisa de manga corta y pantalones cortos.
- Calcetines y zapatos resistentes a productos químicos;
- Guantes impermeables.
- Equipo de protección para la cabeza resistente a productos químicos, para la exposición por encima de la cabeza.

#### PRODUCT INFORMATION

Stalwart 3W is a pre-emergence and post-emergence herbicide for use in corn (field corn, field corn seed, and field corn silage). This product may also be applied to sweet corn, yellow popcorn and grain sorghum as pre-emergence applications prior to crop emergence, or severe adverse crop response may result.

Use this product as a pre-emergence application for control of most annual grasses and broadleaf weeds in the crops described above. This product is also recommended for early post-emergence applications for the control of broadleaf weeds in field corn (pre-emergence applications are recommended in sweet corn, yellow popcorn and grain sorghum). See **Weeds Controlled** section for additional information. This product will not provide consistent control of grasses that have emerged at the time of application.

Stalwart 3W is a mixture of three herbicides, metolachlor, mesotrione, and atrazine with a safener added. It works to control weeds by interfering with the plant's ability to germinate and develop. See **Weeds Controlled** section for additional information and a list of target weed species.

#### USE RATE APPLICATION INFORMATION

Before making applications of this product, determine the soil organic matter content of the field.

- For soils with <3% organic matter content – use 3.0 quarts of this product per acre.
- For soils with >3% organic matter content – use 3.5 quarts of this product per acre.
- Use of this product on soils with >10% soil organic matter is not recommended and may result in poor weed control.

**DO NOT** make applications more than 14 days before planting or to field corn that is taller than 12 inches.

**DO NOT** apply more than 3.5 quarts/A of product per year.

## **USE RESTRICTIONS**

Read all label directions and use instructions before using.

### **• Atrazine Herbicide Rate Limitations**

There may be use rate limitations established in certain states within specific geographical areas for the use of atrazine. These more restrictive and protective requirements must be followed. Consult your state pesticide control agency for additional information. It is a violation of the law and this label to deviate from state use regulations.

- Maximum application rates for atrazine in field corn, field corn seed, field corn silage, sweet corn, and yellow popcorn must be as follows:
  - If no applications of atrazine were made prior to corn emergence, apply a maximum of 2.0 lbs. a.i./A broadcast. If a post-emergence treatment will be required following an earlier herbicide application, the total atrazine applied must not exceed 2.5 lbs. a.i./A per calendar year. When tank mixing or making applications sequentially with atrazine or products containing atrazine to corn, the total pounds of atrazine applied (lbs. a.i./A) must not exceed 2.5 pounds a.i./A per year.
  - Apply a maximum of one 2.0 lbs. a.i./A pre-emergence application on soils that are not highly erodible or on highly erodible soils if at least 30% of the soil is covered with plant residues; or
  - Apply a maximum of one 1.6 lbs. a.i./A pre-emergence application on highly erodible soils if less than 30% of the surface is covered with plant residues, or 2.0 lbs. a.i./A if only applied post-emergence.

**Note:** For calculating total atrazine active ingredient in applications, Stalwart 3W contains 1.71 lbs. a.i. atrazine per gallon.

- **Grazing Restriction:** DO NOT graze or feed forage from treated areas for 45 days following last application, to avoid possible illegal crop residues.
- **Pre-Harvest Interval (PHI):** DO NOT harvest forage, grain, or stover within 60 days after last application of product. Field corn may be treated up to 12 inches tall.
- DO NOT make applications with this product through any type of irrigation system.
- DO NOT make applications under conditions which favor runoff or wind erosion to soil that has been treated with this product or drift to non-target areas.
- To prevent movement to off-site areas due to runoff or wind erosion:
  - When conditions are favorable for wind erosion, avoid treating powdery dry or light sand soils. Allow the soil surface to settle by rainfall or irrigation first under these types of conditions.
  - DO NOT make applications to impervious substrates, such as paved or highly compacted surfaces.
- Unless at least 0.5 inch of rainfall has occurred between the time of application and the first irrigation, DO NOT use tail water from the first flood or furrow irrigation of treated fields to treat non-target crops.
- Where reference is made to weeds partially controlled or suppressed, this can be defined as inconsistent control from good to poor or consistent control at a level below what is typically considered acceptable for commercial weed control.
- Weed control effectiveness may be reduced in dry weather conditions following pre-emergence application of this product. Cultivate the field if weeds develop in conventional tillage corn.
- Observe and follow all precautions and limitations on the label for each tank mix partner product.
- Thoroughly clean sprayer or other application device before using. Dispose of cleaning solution in a responsible manner. **DO NOT** use a sprayer or applicator contaminated with other materials, or crop damage or sprayer clogging of the application device may occur.
- This product will not provide consistent control of most emerged grass weeds.
- **DO NOT** exceed 3.5 qts. of this product per year.
- **DO NOT** make applications of other solo HPPD inhibitor post-emergence herbicides such as mesotrione, tembotrione, or topramezone to areas that have been treated with this product during the same year.
- Severe adverse crop response and corn injury can result if applying this product post-emergence to corn that has emerged and that has received an at-plant application of terbufos insecticide. If this product is applied to emerged corn where an organophosphate insecticide other than terbufos has been applied at planting, temporary corn injury may occur.
- **DO NOT** make applications of this product in a tank mix with any carbamate or organophosphate insecticide to corn that has emerged, or severe corn injury may occur.
- Severe crop injury may result to corn that has emerged with application of any carbamate or organophosphate insecticide applied within 7 days before or 7 days after an application of this product.
- **DO NOT** make applications with this product on any crop other than field corn (for grain, seed, or silage), sweet corn (pre-emergence applications only), yellow popcorn (pre-emergence applications only) or grain sorghum (pre-emergence applications only).
- **DO NOT** make applications with this product in white popcorn or ornamental (Indian) corn or crop injury may occur.
- **DO NOT** contaminate water used for domestic purposes or irrigation water used for crops other than field corn.
- Avoid drift onto adjacent crops and non-target areas.
- Avoid spray overlap, as adverse crop response or crop injury may result.
- **DO NOT** contaminate feed or food with this product.
- **DO NOT** store product near seeds, fertilizers, or foodstuffs.
- Keep all containers of this product tightly closed when not in use.
- **DO NOT MAKE APPLICATIONS OF THIS PRODUCT BY AIR.**

This product will not harm the treated crop if applications are made according to directions and under normal growing conditions. Extended periods of unusually cold and wet or hot and dry weather, insect or plant disease attack, carryover pesticide residues, the use of certain soil-applied systemic insecticides, improperly placed fertilizers or soil insecticides, may weaken crop seedlings during germination and early stages of growth. Using this product under these conditions could result in adverse crop response or crop injury.

### **WEED RESISTANCE MANAGEMENT**

For resistance management, please note that this product contains a Group 5 (atrazine), a Group 15 (metolachlor), and a Group 27 (mesotrione) herbicide. Any weed population may contain plants naturally resistant to Group 5, Group 15, and/or Group 27 herbicides. The resistant individuals may dominate the weed population if these herbicides are used repeatedly in the same fields. Appropriate resistance management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of this product or other Groups 5, 15, or 27 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method such as hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Sipcam Agro at 919-226-1195.

To reduce the risk of weeds developing resistance to HPPD inhibitors, do not make applications of solo post-emergence HPPD inhibitor herbicides (products containing mesotrione or tembotrione) during the same year or on the same field where this product has been applied. A good weed resistance management strategy includes an herbicide spray program that contains two or more modes of action. This product contains three herbicide active ingredients and three modes of action and can be an effective component of a resistance management strategy.

Certain broadleaf weed species have naturally occurring biotypes with resistance to triazines or ALS inhibiting herbicides. No known resistance to this product exists and there are no known instances of cross resistance between this herbicide and other classes of herbicides. If biotypes of triazine or ALS inhibitor-resistant weeds are present in the field and are listed in the **WEEDS CONTROLLED** section of this label, this product should control them.

#### **Integrated Pest (Weed) Management**

Integrate this product into an overall weed pest management strategy whenever the use of an herbicide is required. Practices known to reduce weed development (tillage, crop competition) and herbicide use (weed scouting, proper application timing, banding) should be followed wherever possible. Consult local agricultural and weed authorities for additional IPM strategies established for your area.

#### **Soil Organic Matter**

Before making applications of this product, determine the soil organic matter content of the field. The application use rate is based on percent soil organic matter.

#### **Reduced and No-Till Systems**

This product may be used in reduced and no-till systems. To obtain optimum control, make applications as close to planting as possible. In reduced or no-till systems where weeds are present at application and the corn has not yet emerged, it is recommended to tank mix this product with a burndown herbicide product containing paraquat dichloride or glyphosate.

### **GROUND APPLICATION**

#### **SPRAY EQUIPMENT**

Space spray nozzles uniformly using the same size and type nozzle to provide accurate and uniform application. To avoid drift and produce good coverage, use nozzles that will produce medium to coarse size droplets. Only use 50-mesh or coarser screens in all inline strainer and nozzle screens. Using agitation, maintain proper product dispersion in the tank, and use a pump that can maintain pressure of at least 35 to 40 PSI at the nozzles. If using extended range or drift reduction nozzles, reduced pressure may be used provided that adequate coverage is maintained. Ensure proper and consistent agitation during spraying through duration until spraying is complete – even when there are brief periods of time where spraying has stopped. Stop and run a full agitation before resuming spray if the spray tank is allowed to sit for more than 5 minutes to re-suspend the solution.

#### **Pre-Emergence Applications**

Make pre-emergence applications of this product in a spray volume of 10 to 80 gals/A.

#### **Early Post-Emergence Applications**

For optimum weed control, good weed coverage is essential. For broadcast, over-the-top applications, boom height should be at least 15 inches above the crop canopy, but just high enough to give good, uniform coverage. Make applications in a spray volume of 10 to 30 gals/A. If weed pressure is high and foliage is dense, use a minimum spray volume of 20 gals/A. For post-emergence applications, use flat fan nozzles of 80° or 110° angled forward at 45° for best coverage. Do not use flood jet nozzles or controlled droplet application.

## MANDATORY SPRAY DRIFT MANAGEMENT

### Ground Boom Applications

- **DO NOT** release spray at a height greater than 3 feet above the ground or crop canopy.
- Applicators must select the nozzle and pressure that deliver coarse or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- User must maintain a 15 foot (4.6 meter) in-field downwind buffer (in the direction in which the wind is blowing) from the edge of streams and rivers, as well as high-tide line for all estuarine/marine environments.
- **DO NOT** apply during temperature inversions.

### Boomless Ground Applications

- Applicators must select the nozzle and pressure that deliver coarse or coarser droplets in accordance with American Society of Agricultural & Biological Engineers Standard 572 (ASABE S572).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- User must maintain a 15 foot (4.6 meter) in-field downwind buffer (in the direction in which the wind is blowing) from the edge of streams and rivers, as well as high-tide line for all estuarine/marine environments.
- **DO NOT** apply during temperature inversions.

## SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

### IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### BOOM HEIGHT – Ground Boom

- For ground equipment, the boom should remain level with the crop and have minimal bounce.

### SHIELDED SPRAYERS

- Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

### TEMPERATURE AND HUMIDITY

- When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

### TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

### WIND

Drift potential generally increases with wind speed. AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS. Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

### BOOMLESS GROUND APPLICATIONS:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift. **HANDHELD TECHNOLOGY APPLICATIONS**
- Take precautions to minimize spray drift.

### Sensitive Areas

Apply this product only when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

## ADDITIVES / ADJUVANTS

For applications where an adjuvant will be used, it is recommended to select one that meets the standards of the Chemical Producers and Distributors Association (CPDA) adjuvant certification.

A non-ionic surfactant at 0.25% v/v (1 qt./100 gals.) may be used when this product is applied after field corn has emerged. Using a crop oil concentrate (COC) may result in temporary crop injury. If COC is used as an adjuvant, use at a rate that does not exceed 1% v/v (1 gal./100 gals.) or no more than 1 qt./A. Unless directed for a specific tank mix on this label or as a part of a supplemental label, do not use methylated seed oil (MSO) or nitrogen based adjuvants (AMS or UAN) with this product when applied alone to emerged field corn, or when the application is made as a post-emergence tank mixture with other products. Any of these adjuvants may be used at pre-plant or pre-emergence timing (where the corn plant has not yet emerged) to increase burndown activity on existing weeds. Do not make applications of this product to emerged sweet corn, yellow popcorn or grain sorghum as severe adverse crop response or crop injury may result.

**Tank Mixtures with Glufosinate:** For application of Stalwart 3W tank mixes with registered glufosinate products to emerged field corn, AMS may be added as directed on the glufosinate label. This tank mixture, with registered glufosinate products, should only be applied to LibertyLink® corn hybrids or other corn varieties/cultivars warranted as tolerant to glufosinate. Since the use of other adjuvants may cause severe adverse crop response, AMS should be the only adjuvant added to this tank mix.

**Early Pre-Plant:** Make applications of this product up to 14 days before planting.

**Pre-Emergence Surface:** Do not exceed 3.5 qts. of this product per year. Make applications to the soil surface as a broadcast or banded application.

**Banding Pre-Emergence:** Make applications of this product in a 10-15 inch band after corn planting but before corn emergence.

**Band Applications:** Using row and band width measurements in inches, calculate the amount of product to be applied per acre as follows:

$$\frac{\text{Band width (inches)}}{\text{Row width (inches)}} \times \text{Rate/acre for a broadcast treatment} = \text{Amount needed per acre}$$

**Early Post-Emergence:** Make applications of this product after field corn emergence. Refer to the **Additives** section of this label for adjuvant recommendations. Do not make applications early post-emergence to field corn with liquid fertilizer or severe adverse crop response or crop injury may result. Make applications of this treatment to small broadleaf weeds (no taller than 5 inches) and before the field corn reaches 12 inches in height. Field corn leaf burn may result, but this will not affect later crop growth or yield. Do not make applications of this product to sweet corn or yellow popcorn that has emerged or severe adverse crop response or crop injury may result.

**Emerged Grass Weeds:** This product will not provide consistent control of grass weeds that have emerged. For control of grassweeds that have emerged, a tank mix with another herbicide may be necessary (refer to the **Tank Mixtures** section of this label for additional information).

If applications of metolachlor, s-metolachlor, atrazine (either alone or in combination) have been made prior to an application of this product, limit the early post-emergent application of this product to a total maximum of 2.5 lbs. of active ingredient of atrazine or 3.75 lbs. of metolachlor active per acre, or illegal crop residues may result.

**Sprinkler Irrigation:** **DO NOT** make applications of this product by sprinkler irrigation. After application of this product, a sprinkler irrigation system set to deliver 0.5 to 1 inch of water may be used to incorporate the product. **DO NOT** make applications or incorporate this product with flood irrigation.

## CULTIVATION

A shallow cultivation or rotary hoeing will typically improve weed control if weeds should develop. Cultivate less than half the depth of incorporation if this product was incorporated.

If cultivation becomes necessary because of escaped weeds, compaction, or soil crusting, adjust equipment to run shallow and minimize soil movement. This will decrease the potential of diluting or moving the herbicide away from the weed control zone.

## MIXING PROCEDURES

**Pre-Emergence Applications:** Use either clean water or liquid fertilizers (excluding suspension fertilizers) as carriers for pre-emergence applications. If using fluid fertilizers, a compatibility test must be conducted. See **COMPATIBILITY TEST** section for additional information. Even if this product is determined to be physically compatible with a fluid fertilizer, constant agitation will be necessary to maintain a uniform solution during application.

**Post-Emergence Applications:** When making applications of this product after field corn emergence, use only clean water as the carrier. **DO NOT** make applications to sweet corn or yellow popcorn that has emerged.

### Adding Stalwart 3W to the Spray Tank

The spray tank must be thoroughly rinsed, decontaminated and clean before adding either this product alone or with tank mix partners. Use only clean water, if water is used as the carrier.

**Stalwart 3W Applied Alone:** Add the recommended amount of product to the spray tank when the tank is half full of the carrier, then add the remaining water or fluid fertilizer. Provide agitation during mixing and application so that uniform mixture is maintained.

**Stalwart 3W Applied in Tank Mixtures:** Refer to specific tank mix recommendation sections in this label. Always refer to the tank mix partner label(s) for mixing directions and precautions. **DO NOT** exceed maximum label use rates, or combined total maximum yearly use rates for mesotrione, metolachlor, or atrazine. **DO NOT** mix this product with any product bearing a label prohibition against such mixing. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

If a tank mixture is used, a compatibility test must be conducted. See **COMPATIBILITY TEST** section below for information on conducting a compatibility test.

### COMPATIBILITY TEST

To ensure compatibility of a tank mix partner with this product, a compatibility test should be conducted.

Complete liquid fertilizers or nitrogen solutions (excluding suspension fertilizers) may replace all or part of the water in the spray, as recommended in directions for use. Always conduct compatibility test and make actual applications according to label directions and use recommended carrier. Always check compatibility of liquid fertilizers with pesticide(s) before use because, even within the same analysis, liquid fertilizers vary. Tank mixture incompatibility is more common with mixtures of fertilizers and pesticides.

#### COMPATIBILITY TEST PROCEDURE

(Based on a 25 gal./A spray volume)

1. Add 1.0 pt. of water or fertilizer carrier to each of **two** - 1 quart jars with tight lids. It is important to use the same source of water that will be used in the tank mix and to conduct the test at the same temperature the tank mix will be applied as water and temperature can affect compatibility.
2. Add 1/4 tsp. (or 1.2 mL) of a compatibility agent approved for the intended use to **one of the jars** (1/4 tsp. equals 2.0 pts/100 gals. of spray). Mix by shaking or gently stirring (if shaking place lid on jar).
3. Add the appropriate amount of pesticide(s) based on recommended label rates to **both jars**. If more than one pesticide product will be used, add them separately in the order as described in the **MIXING PROCEDURES** section of this label. Shake or stir gently after each addition to thoroughly mix (if shaking place lid on jar).
4. After all ingredients have been added, place lids on tightly, and invert each jar ten times. Allow the mixtures to stand 15 to 30 minutes. Look for separation, precipitates, gels, heavy oily film on the jar, large flakes, or other signs of incompatibility. Compare the two jars to determine if the compatibility agent is needed. If mixtures separate, but can be easily and readily remixed, the mixture can be sprayed but good agitation must be used. If it is determined the mixtures are incompatible, use the following methods to test for improving compatibility:
  - a. Make a slurry of the dry pesticide(s) in water before addition, or
  - b. Add 1/2 of the compatibility agent to the carrier (fertilizer or water) and the other 1/2 to the emulsifiable concentrate (EC) or flowable pesticide before adding to the mixture. If mixture is still not compatible, do not use the mixture.
5. Dispose of any pesticide wastes in accordance with the Storage and Disposal section in this label.

### TANK MIXTURES

#### Tank Mix Instructions

If the tank mix partner is determined to be compatible, fill the tank half full of the carrier. Begin agitation and maintain throughout mixing and application. Make sure all return lines to the spray tank discharge below the liquid level. Prepare the tank mixture components and add to the tank in the following order:

1. If using a wettable powder or dry flowable formulation, make a slurry with water first and then add it slowly through the screen into the tank. Maintain agitation during this step.
2. If using a flowable formulation, add slowly through screen into the tank. Diluting the flowable with water before adding to the tank may improve mixing and compatibility with dry flowable formulations.
3. Add Stalwart 3W.
4. Add any other tank mix products, adding emulsifiable concentrates last.
5. If an adjuvant will be used, add as the final step. Maintain agitation.
6. Complete filling the spray tank with the carrier and maintain agitation. Make application as soon as possible after spray mixture is prepared. Do not leave mixture in spray tank overnight unattended or without agitation.

#### Cleaning Equipment Post Application

Careful attention must be used when cleaning equipment before spraying a crop other than field corn following applications with this product. Mix the volume of spray solution based on the area of application and mix only as much spray solution as needed.

## Tank and Sprayer Clean Out

1. Use clean water to flush the tank, hoses, boom, and nozzles.
2. Add 1 gal. of household ammonia per 25 gals. of water. Or alternatively, use a commercially available spray tank cleaner.
3. Using pressure washer, clean the inside of the spray tank with this solution. Wash all parts of the tank, including the inside and top surface. If there is not a pressure washer available, fill the sprayer completely with the cleaning solution to provide contact with all internal surfaces of the tank and plumbing. Begin agitation in the sprayer and thoroughly recirculate the solution in the tank for at least 15 minutes. Remove all visible deposits from the spray equipment.
4. Use the cleaning solution to flush the hoses, spray lines, and nozzles for at least 1 minute.
5. Flush dead space areas with water by removing boom end caps, and then replace caps.
6. Dispose of rinsate from the clean-out according to all local State and federal regulations.
7. Repeat the steps 2 to 5 above.
8. After completing the above procedures, remove and clean the nozzles, screens, and strainers separately in the cleaning solution.
9. Completely rinse the spray tank and equipment with clean water.

Make a burndown herbicide application or till the field to destroy emerged or germinating weeds before planting. Immediately after tillage, plant crop into moist soil.

Make applications of this product as directed in this label to control or suppress the weeds listed in the tables below. Tank mixtures may control additional weeds. Always refer to the tank mix partner label(s) for specific use rates, directions and restrictions.

Weed control may be reduced, if a sufficient rainfall is not received within 7 days after application. Apply 0.5 to 1 inch of water, if irrigation is available. Conduct a uniform, shallow cultivation as soon as weeds emerge, if irrigation is not available.

### PRE-EMERGENCE APPLICATIONS: WEED CONTROLLED OR SUPPRESSED

#### BROADLEAF WEEDS

Common Name	Scientific Name	C = Control	S = Suppressed
Amaranth, Palmer	<i>Amaranthus palmeri</i>		C
Amaranth, Powell	<i>Amaranthus powellii</i>		S
Bedstraw, catchweed	<i>Galium aparine</i>		S
Beggarweed, Florida	<i>Desmodium tortuosum</i>		C
Buckwheat, wild	<i>Polygonum convolvulus</i>		C
Buffalobur	<i>Solanum rostratum</i>		C
Carpetweed	<i>Mollugo verticillata</i>		C
Chickweed, common	<i>Stellaria media</i>		C
Cocklebur, common	<i>Xanthium strumarium</i>		S
Deadnettle, purple	<i>Lamium purpureum</i>		C
Devil's claw	<i>Proboscidea louisianica</i>		C
Galinsoga	<i>Galinsoga parviflora</i>		C
Henbit	<i>Lamium amplexicaule</i>		C
Horseweed (marestail)	<i>Conyza canadensis</i>		C
Jimsonweed	<i>Datura stramonium</i>		C
Kochia	<i>Kochia scoparia</i>		C
Lambsquarters, common	<i>Chenopodium album</i>		C
Mallow, Venice	<i>Hibiscus trionum</i>		C
Morningglory, entireleaf	<i>Ipomoea hederacea</i>		S
Morningglory, ivyleaf	<i>Ipomoea hederacea</i>		S
Mustard, wild	<i>Brassica kaber</i>		C
Nightshade, black	<i>Solanum nigrum</i>		C
Nightshade, Eastern black	<i>Solanum pycnanthum</i>		C
Nightshade, hairy	<i>Solanum sarrachoides</i>		C
Pigweed, redroot	<i>Amaranthus retroflexus</i>		C
Pigweed, smooth	<i>Amaranthus hybridus</i>		C
Puncturevine	<i>Tribulus terrestris</i>		S
Purslane, common	<i>Portulaca oleracea</i>		C
Pusley, Florida	<i>Richardia scabra</i>		C
Radish, wild	<i>Raphanus raphanistrum</i>		C
Ragweed, common	<i>Ambrosia artemisiifolia</i>		C
Ragweed, giant	<i>Ambrosia trifida</i>		S
Sesbania, hemp	<i>Sesbania exaltata</i>		C

Shepherd's purse	<i>Capsella bursa-pastoris</i>	C
Sicklepod	<i>Senna obtusifolia</i>	S
Sida, prickly	<i>Sida spinosa</i>	C
Smartweed, ladysthumb	<i>Polygonum persicaria</i>	C
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>	C
Sunflower, common	<i>Helianthus annuus</i>	S
Velvetleaf	<i>Abutilon theophrasti</i>	C
Waterhemp, common	<i>Amaranthus rudis</i>	C
Waterhemp, tall	<i>Amaranthus tuberculatus</i>	C

#### GRASSES

Common Name	Scientific Name	C = Control S = Suppressed
Barnyardgrass	<i>Echinochloa crus-galli</i>	C
Crabgrass	<i>Digitaria spp.</i>	C
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	C
Cupgrass, prairie	<i>Eriochloa contracta</i>	C
Cupgrass, Southwestern	<i>Eriochloa gracilis</i>	C
Cupgrass, woolly	<i>Eriochloa villosa</i>	S
Foxtail, giant	<i>Setaria faberi</i>	C
Foxtail, green	<i>Setaria viridis</i>	C
Foxtail, robust (purple, white)	<i>Setaria spp.</i>	C
Foxtail, yellow	<i>Setaria pumila</i>	C
Goosegrass	<i>Eleusine indica</i>	C
Johnsongrass, seedling	<i>Sorghum halepense</i>	S
Millet, foxtail	<i>Setaria italica</i>	C
Millet, wild proso	<i>Panicum miliaceum</i>	S
Panicum, browntop	<i>Panicum fasciculatum</i>	C
Panicum, fall	<i>Panicum dichotomiflorum</i>	C
Panicum, Texas	<i>Panicum texanum</i>	S
Rice, red	<i>Oryza sativa</i>	C
Sandbur, field	<i>Cenchrus incertus</i>	S
Shattercane	<i>Sorghum bicolor</i>	S
Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>	S
Signalgrass, narrowleaf	<i>Brachiaria piligera</i>	C
Sprangletop, red	<i>Leptochloa filiformis</i>	C
Starbur, bristly	<i>Acanthospermum hispidum</i>	C
Witchgrass	<i>Panicum capillare</i>	C

#### SEDGES

Common Name	Scientific Name	C = Control S = Suppressed
Nutsedge, yellow	<i>Cyperus esculentus</i>	C

#### EARLY POST-EMERGENCE APPLICATIONS\*: WEEDS CONTROLLED OR SUPPRESSED

##### BROADLEAF WEEDS

Common Name	Scientific Name	C = Control S = Suppressed
Amaranth, Palmer	<i>Amaranthus palmeri</i>	C
Amaranth, Powell	<i>Amaranthus powellii</i>	C
Beggarweed, Florida	<i>Desmodium tortuosum</i>	C
Buckwheat, wild	<i>Polygonum convolvulus</i>	C
Buffalobur	<i>Solanum rostratum</i>	C
Carpeweed	<i>Mollugo verticillata</i>	C
Chickweed, common	<i>Stellaria media</i>	C

Cocklebur, common	<i>Xanthium strumarium</i>	C
Dandelion	<i>Taraxacum officinale</i> Weber	S
Deadnettle, purple	<i>Lamium purpureum</i>	C
Devil's claw	<i>Proboscidea louisianica</i>	C
Galinsoga	<i>Galinsoga parviflora</i>	C
Hemp	<i>Cannabis sativa</i> L.	C
Henbit	<i>Lamium amplexicaule</i>	C
Horsenettle	<i>Solanum carolinense</i>	C
Horseweed (marestail)	<i>Conyza canadensis</i>	C
Jimsonweed	<i>Datura stramonium</i>	C
Kochia	<i>Kochia scoparia</i>	C
Lambsquarters, common	<i>Chenopodium album</i>	C
Mallow, Venice	<i>Hibiscus trionum</i>	C
Marestail	<i>Hippuris vulgaris</i> L.	C
Morningglory, entireleaf	<i>Ipomoea hederacea</i>	C
Morningglory, ivyleaf	<i>Ipomoea hederacea</i>	C
Mustard, wild	<i>Brassica kaber</i>	C
Nightshade, black	<i>Solanum nigrum</i>	C
Nightshade, Eastern black	<i>Solanum ptycanthum</i>	C
Nightshade, hairy	<i>Solanum sarrachoides</i>	C
Pigweed, redroot	<i>Amaranthus retroflexus</i>	C
Pigweed, smooth	<i>Amaranthus hybridus</i>	C
Pokeweed	<i>Phytolacca americana</i>	C
Potatoes, volunteer	<i>Solanum spp.</i>	C
Purslane, common	<i>Portulaca oleracea</i>	C
Pusley, Florida	<i>Richardia scabra</i>	C
Radish, wild	<i>Raphanus raphanistrum</i>	C
Ragweed, common	<i>Ambrosia artemisiifolia</i>	C
Ragweed, giant	<i>Ambrosia trifida</i>	C
Sesbania, hemp	<i>Sesbania exaltata</i>	C
Shepherd's purse	<i>Capsella bursa-pastoris</i>	C
Sida, prickly	<i>Sida spinosa</i>	C
Smartweed, ladysthumb	<i>Polygonum persicaria</i>	C
Smartweed, Pennsylvania	<i>Polygonum pensylvanicum</i>	C
Sunflower, common	<i>Helianthus annuus</i>	C
Thistle, Canada	<i>Cirsium arvense</i> (L.) SCOP.	C
Velvetleaf	<i>Abutilon theophrasti</i>	C
Waterhemp, common	<i>Amaranthus rudis</i>	C
Waterhemp, tall	<i>Amaranthus tuberculatus</i>	C

#### GRASSES

Common Name	Scientific Name	C = Control S = Suppressed
Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>	C**

#### SEDGES

Common Name	Scientific Name	C = Control S = Suppressed
Nutsedge, yellow	<i>Cyperus esculentus</i>	S

\*This product will not provide consistent control of most grass weeds that have already emerged.

\*\*Make application prior to weed reaching 2 inches in height.

## ROTATIONAL CROPS

Following application of this product refer to the following rotational crop information:

1. If the crop is lost, field corn, field corn seed, field corn silage, sweet corn, yellow popcorn and grain sorghum (seed treated with Concep® only) may be replanted immediately. Do not make another application of this product.
2. If application of this product is made after June 1, rotating to crops other than corn (all types) or sorghum the following spring may result in crop injury.
3. **DO NOT** rotate to crops other than corn (all types), cotton, small cereal grains, soybeans, sorghum or peanuts the spring following application of this product.
4. For soils having a calcareous surface layer, crop injury may result to soybeans planted the year following application of this product (ex. soils found within the Clarion-Nicollet-Webster soil series in Northern Iowa and Southern Minnesota).
5. **DO NOT** rotate to soybeans for at least 18 months following application of this product if the combined atrazine rate applied was greater than 2.0 lbs. a.i./A, or equivalent band application rate in Dakotas (eastern region), KS, MN (western region), and NE, or soybean crop injury may result.
6. Where rainfall is sparse or erratic or where irrigation is required, in the High Plains and Intermountain areas of the West, use only when corn (all types) or sorghum is to follow field corn, or a crop of untreated corn (all types) or sorghum is to precede other rotational crops.
7. Rotational crop interval is 18 months for all other crops.

---

## CROPS

---

### CORN

---

Stalwart 3W may be used as a pre-emergence application for control of most annual grass and broadleaf weeds in field corn, field corn seed, field silage corn, sweet corn, and yellow popcorn. This product may also be applied as an early post-emergence for the control of broadleaf weeds in field corn, field corn seed, and field silage corn. **DO NOT** make applications of this product to emerged sweet corn or yellow popcorn, or severe adverse crop response or crop injury will result.

See the **Weeds Controlled** tables for a list of weeds controlled or suppressed. This product will not provide consistent control of grasses that have already emerged at the time of application.

#### Use Rate:

- Use 3.0 qts. of this product per acre, if soil organic matter content is less than 3%.
- Use 3.5 qts. of this product per acre, if soil organic matter content is 3% or greater.
- **DO NOT** make applications to field corn taller than 12 inches, or apply more than 3.5 qts./A per year.
- This product is not recommended on soils that have greater than 10% organic matter, or poor weed control may occur.

#### USE RESTRICTIONS:

- **MAXIMUM SINGLE APPLICATION RATE:** 3.5 qts./product/A (1.49 lbs. metolachlor A/A, 0.19 lbs. mesotrione A/A and 1.45 lbs. atrazine A/A)
- **MAXIMUM ANNUAL APPLICATION RATE PER YEAR:** 3.5 qts./product/A (1.49 lbs. metolachlor A/A, 0.19 lbs. mesotrione A/A and 1.45 lbs. atrazine A/A) (Use 3.0 qts. of this product per acre, if soil organic matter content is less than 3%. Use 3.5 qts. of this product per acre, if soil organic matter content is 3% or greater.)
- **MAXIMUM NUMBER OF APPLICATIONS:** 1 (Two applications are allowed if using the split application program)
- For sweet corn, application via mechanically pressurized handguns is prohibited.

#### Product Applications – Alone

**Early Pre-Plant:** Make applications of this product up to 14 days before planting.

**Pre-Emergence Surface:** **DO NOT** exceed 3.5 qts. of this product per year. Make applications to the soil surface as a broadcast or banded application.

**Banded Pre-Emergence:** Make applications of this product in a 10- to 15-inch band after corn planting but before corn has emerged.

**Band Applications:** Using row and band width measurements in inches, calculate the amount of product to be applied per acre as follows:

$$\frac{\text{Band width (inches)}}{\text{Row width (inches)}} \times \text{Rate/acre for a broadcast treatment} = \text{Amount needed per acre}$$

**Early Post-Emergence:** Make applications of this product after field corn has emerged. See the **Additives** section of this label for specific recommendation if using an adjuvant. **DO NOT** apply early post-emergence to field corn in liquid fertilizer or severe adverse crop response or crop injury may result. Make applications of this product to small broadleaf weeds (no greater than 5 inches tall) and before the field corn reaches 12 inches in height. Field corn leaf burn may result, but this will not affect later crop growth or yield. **DO NOT** make applications of this product to sweet corn or yellow popcorn that has emerged or severe adverse crop response or crop injury may result.

This product will not provide consistent control of grass weeds that have emerged. A tank mix with another herbicide may be required for control of emerged weed grasses (Refer to the **Tank Mixtures** section of this label.).

If applications of metolachlor or s-metolachlor, and atrazine (either alone or in combination) have been made prior to an application of this product, limit the amount of Stalwart 3W as an early post-emergent application to no more than a total of 2.5 lbs. of atrazine active ingredient or 3.75 lbs. of metolachlor active ingredient per acre, or illegal crop residues may result.

**Split Application:** Split applications of this product may be made in field corn, field corn seed, and field silage corn. For a split application program, make applications of 1.5 - 2.0 qts./A before crop has emerged, followed by a second application at a rate of 1.25 – 1.75 qts./A as a post-application after corn has emerged. The total amount of this product applied in the split application program must not exceed 3.0 qts./A in soils with less than 3% soil organic matter and must not exceed 3.5 qts./A in soils with greater than 3% organic matter. Reference the **Early Post-Emergence** section above for additional information on post-emergence applications.

**Product Applications - Tank Mixtures Use of Spray Adjuvants in Tank Mixtures**

Used as a pre-emergence herbicide, prior to weed emergence, spray adjuvants make little or no impact on product performance. Where weeds have emerged and the corn has not, an adjuvant may be used with this product applied alone or applied in tank mixture as a burndown application with a tank mix partner burndown herbicide as directed on the individual tank mix partner label(s). Use only adjuvants approved for agricultural crop use. Refer to the **Additives / Adjuvants** section for additional information and directions.

**Reduced Tillage Burndown Combinations** In reduced or no-till corn and prior to crop emergence, tank mixtures of this product with products containing paraquat dichloride or glyphosate will burndown weeds that have emerged. For optimum results, tank mix applications of this product plus paraquat dichloride should be made to weeds that are 1 to 6 inches in height. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Refer to the paraquat dichloride or glyphosate product label for additional information on weeds controlled, directions for use, restrictions and precautions.

**Applications Prior to Crop Emergence: Pre-Emergence Tank Mixtures**

The tank mix products listed in the table below may be used in conventional, reduced, or no-till cropping systems and be applied using the same methods and same application timings as this product unless otherwise directed in the tank mix partner label. Follow all tank mix product labels for directions for use and restrictions. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Be sure to conduct a compatibility test to application. Tank mixtures with 2,4-D may be made, but extreme care should be taken to ensure compatibility before mixing a load or making an application. 2,4-D products, vary greatly with regard to compatibility and should be checked during every application where a water or carrier source, water or carrier temperature, product source, or tank mixture recipe is altered.

**Tank Mix Recommendations for Pre-Emergence Applications with Stalwart 3W**

Tank Mix Recommendation	Scientific Name	Target Use
Atrazine products	Do not exceed 2.0 lbs. a.i./A in total per tank mix, or 2.5 lbs. a.i./A in total per year	Broadleaf and grass weed control improved
Paraquat dichloride	Refer to product label	Burndown of emerged existing weeds
Simazine products	Refer to product label	Broadleaf and grass weed control improved
Glyphosate products	Refer to product label	Burndown of emerged existing weeds.
Lambda-cyhalothrin products	Refer to product label	Insect control (see product label)

**Applications after Crop Emergence: Early Post-Emergence Tank Mix Recommendations**

The tank mix products listed in the table below may be used in conventional, reduced, or no-till systems and may be applied using the same methods and same application timings as this product unless otherwise directed in the tank mix partner label. Follow all tank mix product label directions for use and restrictions.

Conduct a compatibility test before spraying the tank mixture. **DO NOT** make tank mixture applications of this product to sweet corn or yellow popcorn crops that have emerged.

**Tank Mix Recommendations for Early Post-Emergence Weed Control with Stalwart 3W**

Tank Mix*	Rate (Max.)	Objective
Atrazine products	Do not exceed 2.0 lbs. a.i./A in total per tank mix, or 2.5 lbs. a.i./A in total per year	Broadleaf and grass weed control improved
Lambda-cyhalothrin products	Refer to product label	Insect control (see product label)
Nicosulfuron solo products	Refer to product label	Control of grass weeds that have emerged
Rimsulfuron	Refer to product label	Control of grass weeds that have emerged
Glufosinate products (for use only on corn hybrids designated as LibertyLink® or tolerant to glufosinate)	Refer to product label	Control of grass weeds that have emerged

\* Refer to the **Additives** section of this label for recommendations when making applications of this product alone or in tank mixture to field corn that has emerged.

### Stalwart 3W Applications

#### with Glyphosate to Glyphosate Tolerant Corn

Make applications of this product early post-emergence at 2.25 qts./A in tank mix with a solo glyphosate product that is registered for use for over-the-top use in field corn varieties or cultivars warranted as tolerant to glyphosate. To reduce weed competition with the crop, application of this mixture should be targeted to weeds that are 1 to 2 inches. **DO NOT** make applications of this mixture to corn that is higher than 12 inches. If the glyphosate product has an adjuvant included in the formulation (the product label does not call for an adjuvant being added), only spray-grade ammonium sulfate (AMS) at 8.5 lbs./100 gals. should be added to the tank mixture. If the glyphosate product label recommends an adjuvant in addition to AMS, add a non-ionic surfactant (NIS) at 0.25% v/v and AMS to this spray tank mixture. Do not use urea ammonium nitrate (UAN), crop oil concentrate (COC), or methylated seed oil (MSO) type adjuvants in these tank mixtures, or crop injury may result. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Read and follow all directions for use, precautions and restrictions on the glyphosate tank mix partner label.

As an alternative, a pre-emergence application of this product may be made at 2.25 qts./A as part of a two-pass weed control program when followed by a post-emergence application of a glyphosate-containing product in corn varieties or cultivars warranted as tolerant to glyphosate. When this type of application is made, this product will provide reduced competition of the weeds listed in the **Pre-Emergence Applications: Weeds Controlled or Suppressed** table for a period of 30+ days, improving the flexibility in application timing and effectiveness of the glyphosate-based product application. Follow all directions for use, precautions and restrictions on the glyphosate product label.

#### Stalwart 3W Applications in Glufosinate Tolerant Corn

Make early post-emergence applications of this product at 2.25 qts./A in tank mixture with either glufosinate applied over-the-top in LibertyLink® field corn hybrids or other field corn varieties or cultivars warranted as tolerant to glufosinate. To reduce weed competition with the crop, application of this mixture should be targeted to weeds that are 1 to 2 inches in height. **DO NOT** make applications of this mixture to corn that is higher than 12 inches. Ammonium sulfate (AMS) may be used as a spray adjuvant as directed on glufosinate label. AMS should be the only adjuvant used in this tank mix. Do not use urea ammonium nitrate (UAN), crop oil concentrate (COC), non-ionic surfactants (NIS), or methylated seed oil (MSO) type adjuvants in these tank mixes, or crop injury may result. It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Follow all directions for use, precautions, and restrictions on glufosinate product labels.

As an alternative, a pre-emergence application of this product may be made at 2.25 qts./A as part of a two-pass weed control program when followed by a post-emergence application of glufosinate in field corn designated as LibertyLink® field corn hybrids or other field corn varieties or cultivars warranted as tolerant to glufosinate.

When this type of application is made, this product will provide reduced competition of the weeds listed in the **Pre-Emergence Applications: Weeds Controlled or Suppressed** table for a period of 30+ days, improving the flexibility in application timing and effectiveness of the glufosinate application. Follow all directions for use precautions and restrictions on the glufosinate product label being used.

---

## SORGHUM

---

Make a non-incorporated, pre-plant application of this product in sorghum (seed must be treated with Concep®) up to 21 days before planting and up through pre-emergence for weed control. See the **Pre-Emergence Applications: Weeds Controlled or Suppressed** table for a listing of weeds.

Make a broadcast, non-incorporated spray application at 3.0 qts./A starting at 21 days pre-plant and up through planting, but before sorghum has emerged. Making application less than 7 days before the sorghum planting can increase the risk of crop injury, particularly if there is rainfall or irrigation after the application. Symptoms of crop injury include temporary bleaching of young sorghum leaves, or in severe conditions, stunting or partial stand loss. Making the application of this product at greater than 7 days (and no more than 21 days) before the sorghum planting will reduce the risk of adverse crop response.

When making applications of this product before planting, do not incorporate and minimize soil disturbance of the treatment area during planting to minimize the potential for reduced weed control.

Split applications of this product may be made to sorghum as an early pre-plant (7 to 21 day prior to planting), non-incorporated application at 1.5 to 1.75 qts./A followed by a second application of this product made at 1.25 to 1.5 qts./A before the sorghum has emerged. **DO NOT** exceed 3.0 qts./A of product for the split applications.

It is recommended to use a nonionic surfactant (NIS) type adjuvant at 0.25% v/v or a crop oil concentrate (COC) at 1% v/v in the spray solution if weeds are present at the time of application. A spray grade UAN at 2.5% v/v or AMS at 8.5 lbs./100 gallons of spray may also be added in addition to the COC or NIS to the mixture to improve control of weeds that have already emerged. The addition of additives is not recommended, if weeds have not emerged at the time of application.

#### PRODUCT USE RESTRICTIONS:

- **MAXIMUM SINGLE APPLICATION RATE:** 3.0 qts./product/A (1.3 lbs. metolachlor A/A, 0.16 lbs. mesotrione A/A, and 1.25 lbs. atrazine A/A)
- **MAXIMUM ANNUAL APPLICATION RATE PER YEAR:** 3.0 qts./product/A (1.3 lbs. metolachlor A/A, 0.16 lbs. mesotrione A/A, and 1.25 lbs. atrazine A/A)
- **MAXIMUM NUMBER OF APPLICATIONS:** 1 (Two applications are allowed if using the split application program)
- **DO NOT** make applications of this product to sorghum that is grown on sandy soils (sand, sandy loam, or loamy sand).
- **DO NOT** make applications of this product to grain sorghum that has emerged or severe crop injury will result.
- **DO NOT** make applications of this product to sorghum grown for forage, sweet sorghum (sorgo), sudangrass, sorghum-sudangrass hybrids, or dual-purpose sorghum.
- **DO NOT** apply this product to sorghum that is grown south of Interstate 20 (I-20) or east of Highway 277 in the state of Texas.
- **DO NOT** apply atrazine and propazine products to the same sorghum acre.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

### Storage

Keep container tightly closed when not in use. Do not store near seeds, fertilizers, or foodstuffs. Keep away from heat and flame. Ground water contamination may be reduced by diking and flooring of permanent liquid bulk storage sites with an impermeable material.

### Pesticide Disposal

Open dumping is prohibited. Waste resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Rinse spray equipment. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of as described above, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance.

### Container Handling [less than or equal to 5 gallons]

**Non-refillable container.** Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration.

### Container Handling [greater than 5 gallons]

**Non-refillable container.** Do not reuse or refill this container. Offer for recycling if available. Triple rinse container or pressure rinse (or equivalent) promptly after emptying.

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or a mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

Offer for recycling, if available, or puncture and dispose of in a sanitary landfill, or by incineration.

### Container Handling [greater than 5 gallons]

**Refillable container.** Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank.

Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

**DO NOT USE CONTAINERS FOR THE STORAGE OF FOOD, FEED, OR DRINKING WATER!**

## ALMACENAMIENTO Y DESECHO

No contamine agua, comida ni forrajes mediante el almacenamiento y desecho.

### Almacenamiento

Mantenga el envase bien cerrado cuando no se utilice. No lo almacene cerca de semillas, fertilizantes o alimentos. Manténgalo alejado del calor y de las llamas. La contaminación de las aguas subterráneas se puede reducir construyendo diques y cubriendo los suelos de los lugares de almacenamiento permanente de materiales líquidos a granel con un material impermeable.

### Desecho de pesticidas

El vertido a cielo abierto está prohibido. Los residuos resultantes del uso de este producto deben eliminarse in situ o en un punto de eliminación de residuos autorizado. Enjuague los equipos de rociado.

Los residuos de pesticidas son muy peligrosos. La disposición inadecuada del pesticida sobrante, mezcla de aerosol o aguas de enjuague de contenedores, constituye una violación de la ley federal. Si estos residuos no se pueden eliminar como se ha descrito anteriormente, contacte a la agencia para el manejo de pesticidas o control ambiental de su estado o al representante para el Manejo de Residuos Peligrosos de la Oficina Regional de la EPA más cercana para obtener indicaciones del proceso a seguir para su disposición.

### Manipulación del envase [con una capacidad inferior o igual a 5 galones]

Envase no rellenable. No reutilice ni rellene este envase. Ofrézcalo para ser reciclado en caso de ser posible. Enjuague tres veces el envase (o equivalente) inmediatamente después de vaciarlo.

Enjuague tres veces de la siguiente manera: Vacíe el contenido restante en el equipo de aplicación o en un tanque de mezcla. Drene durante 10 segundos después de que el flujo comience a gotear. Llene el envase hasta 1/4 de su capacidad con agua y vuelva a taparlo. Agite durante 10 segundos. Vierta el agua de enjuague en el equipo de aplicación o en un tanque de mezcla, o almacénela para usarla y desecharla más adelante. Drene durante 10 segundos después de que el flujo comience a gotear. Repita este procedimiento dos veces más. Ofrezca el envase para ser reciclado (en caso de ser posible), o perforelo y deséchelo en un relleno sanitario apropiado, o destrúyalo mediante incineración.

### Manipulación del envase [con una capacidad superior a 5 galones]

Envase no rellenable. No reutilice ni rellene este envase. Ofrézcalo para ser reciclado en caso de ser posible. Enjuague tres veces el envase o enjuáguelo a presión (o equivalente) inmediatamente después de vaciarlo.

Enjuague tres veces de la siguiente manera: Vacíe el contenido restante en el equipo de aplicación o en un tanque de mezcla. Llene el envase hasta 1/4 de su capacidad con agua. Reemplace y ajuste los cierres. Incline el envase sobre un costado y gírelo hacia adelante y hacia atrás, asegurándose de dar al menos una vuelta completa, durante 30 segundos. Coloque el envase sobre un extremo e inclínelo hacia adelante y hacia atrás varias veces. Gire el envase sobre el otro extremo e inclínelo hacia adelante y hacia atrás varias veces. Vacíe el agua de enjuague en el equipo de aplicación o en un tanque de mezcla, o almacénela para usarla o desecharla más adelante. Repita este procedimiento dos veces más.

Enjuague a presión de la siguiente manera: Vacíe el contenido restante en el equipo de aplicación o en un tanque de mezcla y continúe drenando durante 10 segundos después de que el flujo comience a gotear. Sostenga el envase boca abajo sobre el equipo de aplicación o un tanque de mezcla, o recoja el agua de enjuague para usarla o desecharla más adelante. Inserte la boquilla de enjuague a presión en el costado del envase y enjuague a aproximadamente 40 PSI durante al menos 30 segundos. Drene durante 10 segundos después de que el flujo comience a gotear. Ofrezca el envase para ser reciclado (en caso de ser posible), o perforelo y deséchelo en un relleno sanitario apropiado, o destrúyalo mediante incineración.

### Manipulación del envase [con una capacidad superior a 5 galones]

Envase recargable. Recargue este envase únicamente con pesticidas. No reutilice este envase para ningún otro fin. La limpieza del envase antes de desecharlo definitivamente es responsabilidad de la persona que se encarga de desechar el envase. La limpieza del envase antes de recargarlo es responsabilidad de quien recarga el envase. Para limpiar el envase antes de eliminarlo definitivamente, vacíe el contenido restante en el equipo de aplicación o en un tanque de mezcla. Llene el envase hasta un 10 % de su capacidad con agua. Agítelo energicamente o haga circular el agua de nuevo con la ayuda de la bomba por 2 minutos. Vierta o bombee el agua de enjuague en el equipo de aplicación o en el sistema de recolección de agua de enjuague. Repita este procedimiento de enjuague dos veces más.

A continuación, ofrezca el envase para ser reciclado (en caso de ser posible) o perforelo y deséchelo en un relleno sanitario apropiado, o destrúyalo mediante incineración u otros procedimientos aprobados por las autoridades estatales y locales.

**NO UTILICE LOS ENVASES PARA ALMACENAR ALIMENTOS, PIENSOS O AGUA POTABLE.**

## WARRANTY AND LIMITATIONS OF DAMAGES

**Conditions of Sale:** To the extent consistent with applicable law, Sipcam Agro USA, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under conditions not reasonably foreseeable to Sipcam Agro USA, Inc. Sipcam Agro USA, Inc. disclaims all other warranties, express or implied. To the extent consistent with applicable law, Sipcam Agro USA, Inc. shall not be liable for consequential, special, or indirect damages resulting from the use or handling of this product, and Sipcam Agro USA, Inc.'s sole liability and buyer's and user's exclusive remedy shall be limited to the refund of the purchase price. Buyer and user acknowledge and assume all risks and liability resulting from handling, storage, and use of this product. Sipcam Agro USA, Inc. does not authorize any agent or representative to make any other warranty, guarantee or representation concerning this product.

Concept® is a trademark of Syngenta Group Company

LibertyLink® is a trademark of Bayer CropScience

# RESTRICTED USE PESTICIDE

## DUE TO GROUND AND SURFACE WATER CONCERNS

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION, AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION. THIS PRODUCT IS A RESTRICTED-USE HERBICIDE DUE TO GROUND AND SURFACE WATER CONCERNS. USERS MUST READ AND FOLLOW ALL PRECAUTIONARY STATEMENTS AND INSTRUCTIONS FOR USE IN ORDER TO MINIMIZE POTENTIAL FOR ATRAZINE TO REACH GROUND AND SURFACE WATER

# Stalwart® 3W

For pre-emergence and post-emergence control of grass and broadleaf weeds in Corn (field, seed, silage, sweet, and yellow popcorn), and Grain Sorghum

### ACTIVE INGREDIENTS:

Metolachlor .....	19.00%
Atrazine .....	18.60%
Mesotrione .....	2.44%

### OTHER INGREDIENTS:

.....	59.96%
-------	--------

TOTAL: .....100.00%

Equivalent to 1.70 lbs.ai./gal. metolachlor; 0.218 lbs.ai./gal. mesotrione; and 1.66 lbs.ai./gal. atrazine.

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

### FIRST AID

#### IF IN EYES:

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.
- Call a poison control center or doctor for treatment advice.

#### IF SWALLOWED:

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

#### IF ON SKIN OR CLOTHING:

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

#### IF INHALED:

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
- Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

Emergency Phone Numbers

Poison Control: 1-800-222-1222  
CHEMTREC: 1-800-424-9300 (transportation and spills)

See additional Precautionary Statements and Directions for Use inside the booklet.

Manufactured for:  
Sipcam Agro USA, Inc.  
2525 Meridian Parkway  
Durham, NC 27713

For Spanish translation, see inside the booklet. (Para la traducción al español, consulte el interior del folleto.)

EPA Reg. No.: 60063-64

EPA Est. No.: 60063-GA-1 (Lot number begins with VL)

EPA Est. No.: 62171-MS-1 (Lot number begins with OI)

EPA Est. No.: 70989-MO-1 (Lot number begins with ST)

EPA Est. No.: 72344-MO-4 (Lot number begins with TR)

EPA 20230120 (9/25)



**SIPCAM AGRO**  
USA, INC.

Not for sale, use, or distribution in Nassau County or Suffolk County, New York.

**READ THE LABEL  
CAREFULLY  
BEFORE OPENING  
THE CONTAINER**

HERBICIDE

Metolachlor	Group	15	Herbicide
Mesotrione	Group	27	Herbicide
Atrazine	Group	5	Herbicide