



DIRECTIONS FOR USE

- Product should be used during ideal growing conditions to ensure effective uptake. Avoid extreme weather conditions such as heat and frost events. Spraying in the early morning or late afternoon can assist in minimising risk of rapid drying on the leaf due to evaporation. Where local experience of successful use is not available, we recommend a small scale jar test and test application prior to wide spread use. If any issues are observed do not continue with the application.
- It is recommended to add product to the tank mix after all other products have been mixed well in the tank, including most of the water.

COMPATIBILITY

Compatible with most common fertilisers and crop protection products. It is recommended to perform a jar test before mixing.

SAFETY DATA SHEET

For further information, refer to the Safety Data Sheet (SDS) which is available from the supplier or from our web site, www.sipcam.com.au

G.H.S STATEMENTS - DANGER

Hazard statements

H318 - Causes serious eye damage

H335 - May cause respiratory irritation

Precautionary Statements – Prevention

Avoid breathing dust/fume/gas/mist/vapors/spray.

Use only outdoors or in a well-ventilated area. Wear eye/face protection. Wash hands and face thoroughly after handling.

Precautionary Statements – Response

IF exposed or concerned: IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing.

Immediately call a POISON CENTER or doctor/physician. IF INHALED: Remove person to fresh air and keep comfortable for breathing. Call a POISON CENTER or doctor/physician if you feel unwell.

Precautionary Statements – Storage

Store in a well-ventilated place. Keep container tightly closed. Store locked up.

Precautionary Statements – Disposal

Dispose of contents/container in accordance with local, regional, national, and international regulations as applicable.




Sipcam Pacific
Australia Pty Ltd
Level 1,
191 Malop Street
Geelong
VICTORIA 3220

1000 L

Macrosol K-Carb



K

Potassium (K): 400 g/L as potassium carbonate.

A high analysis potassium carbonate solution, free of any chloride, specifically designed for fertigation and furrow injection applications.

CROPS	RATE	COMMENTS
BROADACRE CROPS	3-10 L/ha	Apply furrow irrigation.
HORTICULTURAL CROPS AND PASTURE	5-30 L/ha	Apply via fertigation as required.

CAUTION: Applications in adverse spraying conditions or in conjunction with some crop protection products may result in reductions in efficacy or phytotoxicity which may not be acceptable to growers. For most effective results apply early morning or late afternoon.

This product is NOT a Dangerous Good under the Australian Code for the Transport of Dangerous Goods by Road and Rail.

Specialist Advice In Emergency Only
1800 033 111 All Hours - Australia Wide

WARNING STATEMENTS

Do not swallow. The spray from this product may act as an irritant. Avoid inhalation and contact with the eyes and skin.

Batch No:

Date of Manufacture:

THIS LABEL COMPLIES WITH THE NATIONAL CODE OF PRACTICE FOR FERTILIZER DESCRIPTION AND LABELLING

NOTICE TO BUYER This product must be used in accordance with the directions for use. Where this product is not a good of a kind ordinarily acquired for personal, domestic or household use or consumption under the Australian Consumer Law, Sipcam's liability for any direct, indirect or consequential loss which you incur, including, without limitation, loss or damage to crop, loss of equipment, property damage, personal injury or death is limited, at Sipcam's option, to the replacement of the product or the supply of an equivalent product. By opening this package, you agree to be bound by these terms. If you do not agree to these terms, please return the entire unopened package intact to Sipcam or the place in which you purchased this product for a full refund.